



Queensland University of Technology
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

Pritchard, Nicola, Edwards, Katie, Srinivasan, Sangeetha, Dehghani, Cirous, Vagenas, Dimitrios, Russell, Anthony W., Malik, Rayaz A., & Efron, Nathan
(2013)

Corneal nerve fibre length is reduced in diabetic individuals who develop 3-year incident diabetic neuropathy. In
International Diabetes Foundation 2013, 2-6 December 2013, Melbourne, Vic.

This file was downloaded from: <https://eprints.qut.edu.au/75645/>

Notice: *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

<http://conference.idf.org/IDF2013/CM.NET.WebUI/CM.NET.WEBUI.SCP2/SCP2functiondetail.asp?0000-0000-0000-00000000000002&sesID=05000000-0000-0000-0000-0000000000967&absID=07000000-0000-0000-0000-000000005448>



search abstracts

OP-0605

Authors

Sessions

Bookmarks

Latest Search Results

All Abstracts (1957)	IDF Award Lecture (8)	Meet-the-Expert (25)	Symposium (304)	Teaching Lecture (14)	Open Forum (22)
Speaker's Corner (4)	Debate (10)	Workshop (21)	Q&A (3)	Poster Discussion (377)	Oral Presentation (227)
Named Lecture (1)	President's Address (1)				Poster (940)

Melbourne 2013

Neuropathy, the foot and the brain

Abstract: OP-0605

Corneal nerve fibre length is reduced in diabetic individuals who develop 3-year incident diabetic neuropathy

N. Pritchard Pritchard¹, K. Edwards Edwards¹, S. Srinivasan Srinivasan¹, C. Dehghani Dehghani¹, D. Vagenas Vagenas¹, A. Russell Russell², R. Malik Malik³, N. Efron Efron¹

¹Queensland University of Technology, Institute of Health and Biomedical Innovation, Brisbane, Australia

²Princess Alexandra Hospital and University of Queensland, School of Medicine, Brisbane, Australia

³University of Manchester and Central Manchester Foundation Trust, Division of Cardiovascular Medicine, Manchester, United Kingdom

N. Pritchard¹, K. Edwards¹, S. Srinivasan¹, C. Dehghani¹, D. Vagenas¹, A. Russell², R. Malik³, N. Efron¹.

¹Queensland University of Technology, Institute of Health and Biomedical Innovation, Brisbane, Australia.

²Princess Alexandra Hospital and University of Queensland, School of Medicine, Brisbane, Australia.

³University of Manchester and Central Manchester Foundation Trust, Division of Cardiovascular Medicine, Manchester, United Kingdom.

Aims:

Depletion of corneal nerves has been investigated as a potential marker and/or predictor of neuropathy. To determine the clinical measures that predict 3-year incidence of diabetic neuropathy, baseline clinical parameters of individuals with type 1 diabetes who developed neuropathy were compared to those who did not develop neuropathy over a 3-year period.

Methods:

Sixty-four participants with diabetes, but without baseline neuropathy in the LANDMark Study Cohort have completed 3 years of longitudinal follow-up. The baseline characteristics of those who developed neuropathy were compared to those that did not. Neuropathy was defined, by the Toronto criteria, as abnormal nerve conduction, and a sign (neuropathy disability score ≥ 3) or symptom (diabetic neuropathy symptom score ≥ 1) of neuropathy.

Results:

Diabetic neuropathy developed in 7 (11%) participants over a mean follow-up of 3.0 (range 2.9 to 3.2) years. Baseline corneal nerve fibre length (CNFL) was significantly lower in individuals who developed neuropathy over 3 years (13.3 ± 3.7 mm/mm²) compared to those that did not (17.4 ± 5.0 mm/mm²) ($P = 0.036$). Reduced peroneal CV (40.6 ± 4.1 vs. 46.7 ± 4.6 , $p = 0.003$), and cold sensation (23.5 ± 5.9 vs. 27.2 ± 5.0 , $p = 0.022$) and increased vibration threshold (20.5 ± 14.4 vs. 11.9 ± 17.6 , $p = 0.034$) were also associated with development of diabetic neuropathy. The odds ratio of having reduced CNFL in those who developed neuropathy vs. those who did not develop neuropathy was 0.871 (95% CI 0.825-0.920) compared to peroneal CV 0.953 (95% CI 0.936-0.970), cold sensation 0.921 (95% CI 0.892-0.950) and vibration thresholds 0.936 (95% CI 0.891-0.982).

Conclusion:

Corneal confocal microscopy examination may be an appropriate adjunct to diabetic neuropathy screening in a clinical setting. Non-invasive corneal evaluation as part of routine eye examination may assist in early detection of peripheral neuropathy and be a useful test in predicting individuals who develop neuropathy.

No conflict of interest